

## · 专家共识 ·

# 中国髋、膝关节置换术围术期抗纤溶药序贯抗凝血药应用方案的专家共识\*

国家卫生计生委公益性行业科研专项《关节置换术安全性与效果评价》项目组

岳辰<sup>1</sup> 周宗科<sup>1</sup> 裴福兴<sup>1\*\*</sup> 翁习生<sup>2\*\*</sup> 邱贵兴<sup>2\*\*</sup> 阮长耿<sup>3\*\*</sup>

(1.四川大学华西医院骨科,成都 610041;2.中国医学科学院北京协和医院骨科,北京 100730;3.江苏省血液研究所,苏州 215006)

髋、膝关节置换术常可伴随大量失血。根据文献报道,髋、膝关节置换术围术期总失血量多在1000 ml以上,输血率高达30%~60%<sup>[1,2]</sup>。大量失血可增加患者的围术期风险和经济负担<sup>[3]</sup>。髋、膝关节置换术围术期失血除手术切口直接出血外,由手术创伤引起的纤溶反应增强所致的失血约占总失血量的60%<sup>[4]</sup>。而且,膝关节置换术中应用止血带引起的组织缺血再灌注损伤可进一步增强纤溶反应<sup>[5]</sup>,增加出血量。

氨甲环酸(tranexamic acid, TXA)是一种抗纤溶药,其与纤溶酶原的赖氨酸结合位点具有高亲和性,可封闭纤溶酶原的赖氨酸结合位点,使纤溶酶原失去与纤维蛋白结合的能力,导致纤溶活性降低,从而发挥止血作用<sup>[6]</sup>。目前,大量研究均已证实氨甲环酸能有效减少髋、膝关节置换术围术期的失血量并降低输血率,且不增加术后静脉血栓栓塞症的发生风险<sup>[5-8]</sup>。

髋、膝关节置换术患者是静脉血栓栓塞症的高发人群,应用抗凝血药物能有效降低静脉血栓栓塞症的发生率。为了在髋、膝关节置换术围术期更好地平衡抗纤溶药与抗凝血药的应用,既可减少患者的出血量、降低输血率,又不增加患者发生静脉血栓栓塞症的风险,保障医疗安全。国家卫生计生委公益性行业科研专项《关节置换术安全性与效果评价》项目组(项目编号:201302007)和《中华骨与关节外科杂志》编辑部邀请国内专家,复习国内外27篇meta分析和260多篇论著,结合项目组26家大型医院数据库和50家推广医院数据库共13300例髋、膝关节置换术病例中8426例氨甲环酸应用经验以及全国12场氨甲环酸临床应用区域会议征求意见结果,遵循循证医学原则,达成髋、膝关节置换术围术期抗纤溶药序贯抗凝血药应用的专家共识,供广大骨科医师

在临床工作中参考应用。但在应用氨甲环酸前应结合患者的全身情况,参照氨甲环酸药物说明书或《中国药典》,遇有不良反应及时处理。

## 1 髋关节置换术围术期的氨甲环酸应用

### 1.1 静脉应用

11篇meta分析<sup>[8-18]</sup>及19篇前瞻性随机对照研究<sup>[19-37]</sup>报道氨甲环酸给药方式主要为单次静脉滴注或二次间隔静脉滴注,二次给药间隔时间为3 h。单次给药剂量为15~20 mg/kg或总量1 g;二次间隔给药剂量为每次10~20 mg/kg或每次总量1 g。

推荐:①单次给药法:髋关节置换术切开皮肤前5~10 min氨甲环酸15~20 mg/kg或总量1 g静脉滴注完毕;②二次间隔给药法:首次给药同单次给药法,3 h后根据引流情况再次给药,剂量同前。

### 1.2 局部应用

研究表明,氨甲环酸局部应用能够提高局部药物浓度,减少全身吸收<sup>[38]</sup>。1篇meta分析<sup>[39]</sup>及4篇前瞻性随机对照研究<sup>[38,40-42]</sup>报道氨甲环酸2~3 g局部应用可以有效减少出血、降低输血率。目前,有关氨甲环酸的局部应用尚无统一标准,特别是对于术后是否放置引流管及引流管夹闭后何时开放仍存在争议,各报道中术后引流管夹闭时间为30 min~2 h不等。因此,氨甲环酸在髋关节置换术中局部应用的具体方法及术后引流管夹闭时间有待进一步研究。

推荐:氨甲环酸在髋关节置换术中局部应用的推荐剂量为2~3 g。

### 1.3 静脉和局部联合应用

研究报道,氨甲环酸在髋关节置换术围术期静脉滴注联合局部应用相比单纯静脉滴注或局部应用能更有效减少出血、降低输血率<sup>[43]</sup>。具体方法为髋关

\*基金项目:国家卫生计生委公益性行业科研专项(项目编号:201302007)

\*\*通信作者:裴福兴,E-mail:peifuxing@vip.163.com;翁习生,E-mail:xshweng@medmail.com.cn;邱贵兴,E-mail:qguixing@126.com;  
阮长耿,E-mail:changgengruan@hotmail.com

节置换术切开皮肤前5~10 min氨甲环酸15~20 mg/kg静脉滴注完毕,同时关闭切口前以总量1~2 g氨甲环酸局部应用。

推荐:髋关节置换术切开皮肤前5~10 min氨甲环酸15~20 mg/kg静脉滴注完毕,同时关闭切口前氨甲环酸1~2 g局部应用。

## 2 髋关节置换术围术期的氨甲环酸应用

### 2.1 静脉应用

13篇meta分析<sup>[7,9,44-54]</sup>及16篇前瞻随机对照研究<sup>[5,19,55-68]</sup>报道,氨甲环酸给药方式主要为单次静脉滴注或二次间隔静脉滴注,二次给药间隔为3 h。单次给药时间应在手术开始前(不用止血带者)或松止血带前5~10 min,剂量为10~20 mg/kg或总量1 g;二次给药时间为首次给药后3 h再次给药,剂量为每次10~20 mg/kg或每次总量1 g。

推荐:①单次给药法:膝关节置换术切开皮肤前(不用止血带者)或松止血带前5~10 min氨甲环酸10~20 mg/kg或1 g静脉滴注完毕;②二次间隔给药法:首次给药同单次给药法,3 h后根据引流情况再次给药,剂量相同。

### 2.2 局部应用

4篇meta分析<sup>[69-72]</sup>及12篇前瞻随机对照研究<sup>[73-84]</sup>报道氨甲环酸局部应用的最低有效剂量 $\geq 2 \text{ g}$ <sup>[70]</sup>、最低有效浓度 $\geq 20 \text{ mg/ml}$ <sup>[71]</sup>,大剂量( $\geq 2 \text{ g}$ )和高浓度( $\geq 20 \text{ mg/ml}$ )氨甲环酸局部应用能有效减少膝关节置换术围术期出血、降低输血率。局部应用方法为关闭切口前关节腔灌注,或关闭切口后通过引流管逆行注入,或通过注射器关节腔内注射。各报道中术后引流管夹闭时间为30 min~2 h不等,仍存在争议,有待进一步研究。

推荐:氨甲环酸在膝关节置换术中的局部应用应在关闭切口前后,局部应用的剂量 $\geq 2 \text{ g}$ 或浓度 $\geq$

20 mg/ml。

### 2.3 静脉和局部联合应用

联合给药方法为松开止血带5~10 min前氨甲环酸15~20 mg/kg或1 g静脉滴注,同时关闭切口前氨甲环酸1~2 g局部注入。联合用药能有效减少膝关节置换术围术期出血、降低输血率<sup>[85]</sup>。

推荐:膝关节置换术切开皮肤前(不用止血带者)或松止血带前5~10 min氨甲环酸15~20 mg/kg或1 g静脉滴注完毕,同时关闭切口前氨甲环酸1~2 g局部应用。

## 3 髋、膝关节置换术围术期抗纤溶药序贯抗凝血药应用

髋、膝关节置换术围术期应用抗纤溶药氨甲环酸后序贯应用抗凝血药,既能减少出血,又不增加静脉血栓栓塞症发生风险。氨甲环酸的止血效果与其应用剂量和应用次数有关,但随着剂量或次数的增加,静脉血栓栓塞症的发生风险也可能增大。理论上认为,抗凝血药物在术后应用越早、持续时间越长,患者发生静脉血栓栓塞症的风险越小,但发生出血的风险增大。为了达到抗纤溶药和抗凝血药的平衡,应在髋、膝关节置换术围术期应用氨甲环酸6 h后根据引流量的变化,选择抗凝血药应用时间。大部分患者术后6~12 h内伤口出血趋于停止,如引流管无明显出血或引流管血清已分离则表明伤口出血趋于停止,应在12 h内应用抗凝血药;若个别患者术后12 h仍有明显出血,可延后应用抗凝血药。

髋、膝关节置换术后抗凝血药物预防持续时间应根据《中国骨科大手术静脉血栓栓塞症预防指南》,推荐预防时间最短为10 d,可延长至11~35 d<sup>[86]</sup>。在应用时应注意抗凝血药物的有效性和安全性,当患者出现凝血功能异常或出血事件时,应综合评价出血与血栓的风险,及时调整药物剂量或停用。

本共识制定专家和项目组专家共54人,各位专家在共识制定过程中贡献了自己的宝贵经验和建议,才使得共识达到了更高的学术水平和应用价值,特向各位专家致以衷心地感谢!

(按姓氏笔画为序)

马信龙	王万春	王兆铖	王坤正	左建林	田伟	王浩洋	沈彬	张先龙	张伟	张克
张英泽	史占军	包倪荣	邢欣	朱庆生	朱振安	朱锦宇	刘军	阮长耿	孙俊刚	孙维
严广斌	严世贵	李庭	杨柳	吴俣	吴海山	邱贵兴	余楠生	张耀南	尚希福	易群
岳辰	周宗科	赵永强	赵建宁	胡永成	胡豫	姜保国	胥伯勇	袁宏	钱齐荣	徐海东
翁习生	高忠礼	高鹏	梅恒	曹力	堪武生	谢锦伟	裴福兴	廖威明	薛庆云	

## 参 考 文 献

- [1] Bierbaum BE, Callaghan JJ, Galante JO, et al. An analysis of blood management in patients having a total hip or knee arthroplasty. *J Bone Joint Surg Am*, 1999, 81(1): 2-10.
- [2] Lemaire R. Strategies for blood management in orthopaedic and trauma surgery. *J Bone Joint Surg Br*, 2008, 90(9): 1128-1136.
- [3] Vamvakas EC, Blajchman MA. Transfusion-related mortality: the ongoing risks of allogeneic blood transfusion and the available strategies for their prevention. *Blood*, 2009, 113(15): 3406-3417.
- [4] Liu X, Zhang X, Chen Y, et al. Hidden blood loss after total hip arthroplasty. *J Arthroplasty*, 2011, 26(7): 1100-1105. e1.
- [5] Engel JM, Hohaus T, Ruwoldt R, et al. Regional hemostatic status and blood requirements after total knee arthroplasty with and without tranexamic acid or aprotinin. *Anesth Analg*, 2011, 92(3): 775-780.
- [6] Hoylaerts M, Lijnen HR, Collen D. Studies on the mechanism of antifibrinolytic action of tranexamic acid. *Biochim Biophys Acta*, 1981, 673(1): 75-85.
- [7] Poeran J, Rasul R, Suzuki S, et al. Tranexamic acid use and postoperative outcomes in patients undergoing total hip or knee arthroplasty in the United States: retrospective analysis of effectiveness and safety. *BMJ*, 2014, 349: g4829.
- [8] Gandhi R, Evans HM, Mahomed SR, et al. Tranexamic acid and the reduction of blood loss in total knee and hip arthroplasty: a meta-analysis. *BMC Res Notes*, 2013, 6: 184.
- [9] Ho KM, Ismail H. Use of intravenous tranexamic acid to reduce allogeneic blood transfusion in total hip and knee arthroplasty: a meta-analysis. *Anaesth Intensive Care*, 2003, 31(5): 529-537.
- [10] Sukeik M, Alshryda S, Haddad FS, et al. Systematic review and meta-analysis of the use of tranexamic acid in total hip replacement. *J Bone Joint Surg Br*, 2011, 93(1): 39-46.
- [11] Pinzón-Florez CE, VélezCañas KM, DíazQuijano DM. Efficiency of tranexamic acid in perioperative blood loss in hip arthroplasty: a systematic literature review and meta-analysis. *Rev Esp Anestesiol Reanim*, 2015, 62(5): 253-264.
- [12] Zhou XD, Tao LJ, Li J, et al. Do we really need tranexamic acid in total hip arthroplasty? A meta-analysis of nineteen randomized controlled trials. *Arch Orthop Trauma Surg*, 2013, 133(7): 1017-1027.
- [13] Gill JB, Rosenstein A. The use of antifibrinolytic agents in total hip arthroplasty: a meta-analysis. *J Arthroplasty*, 2006, 21(6): 869-873.
- [14] 岳辰, 康鹏德, 沈彬, 等. 氨甲环酸用于首次髋关节置换术的系统评价和Meta分析. 中国矫形外科杂志, 2013, 21(12): 1167-1172.
- [15] 周磊, 李涛, 翁习生, 等. 氨甲环酸在全髋关节置换术围手术期疗效与安全性的meta分析. 中国骨与关节外科, 2013, 6(5): 421-426.
- [16] 尹勇, 马广文, 黄斐, 等. 氨甲环酸减少全髋关节置换失血量的Meta分析. 中国组织工程研究, 2014, 18(17): 2752-2757.
- [17] 刘丙根, 庞清江. 氨甲环酸用于全髋关节置换有效性与安全性的Meta分析. 中国组织工程研究, 2014, 18(35): 5699-5706.
- [18] 付鑫, 李稚君, 马信龙, 等. 全髋关节置换术使用氨甲环酸有效性及安全性的Meta分析. 中华关节外科杂志(电子版), 2014, 8(1): 84-90.
- [19] Ido K, Neo M, Asada Y, et al. Reduction of blood loss using tranexamic acid in total knee and hip arthroplasties. *Arch Orthop Trauma Surg*, 2000, 120(9): 518-520.
- [20] Benoni G, Fredin H, Knebel R, et al. Blood conservation with tranexamic acid in total hip arthroplasty: a randomized, double-blind study in 40 primary operations. *Acta Orthop Scand*, 2001, 72(5): 442-448.
- [21] Benoni G, Lethagen S, Nilsson P, et al. Tranexamic acid, given at the end of the operation, does not reduce postoperative blood loss in hip arthroplasty. *Acta Orthop Scand*, 2000, 71(3): 250-254.
- [22] Ekbäck G, Axelsson K, Ryttberg L, et al. Tranexamic acid reduces blood loss in total hip replacement surgery. *Anesth Analg*, 2000, 91(5): 1124-1130.
- [23] Husted H, Blønd L, Sonne-Holm S, et al. Tranexamic acid reduces blood loss and blood transfusions in primary total hip arthroplasty: a prospective randomized double-blind study in 40 patients. *Acta Orthop Scand*, 2003, 74(6): 665-669.
- [24] Lemay E, Guay J, Côté C, et al. Tranexamic acid reduces the need for allogenic red blood cell transfusions in patients undergoing total hip replacement. *Can J Anaesth*, 2004, 51(1): 31-37.
- [25] Garneti N, Field J. Bone bleeding during total hip arthroplasty after administration of tranexamic acid. *J Arthroplasty*, 2004, 19(4): 488-492.
- [26] Yamasaki S, Masuhara K, Fuji T. Tranexamic acid reduces blood loss after cementless total hip arthroplasty- prospective randomized study in 40 cases. *Int Orthop*, 2004, 28(2): 69-73.
- [27] Johansson T, Pettersson LG, Lisander B. Tranexamic acid in total hip arthroplasty saves blood and money: a randomized, double-blind study in 100 patients. *Acta Orthop*, 2005, 76(3): 314-319.
- [28] Niskanen RO, Korkala OL. Tranexamic acid reduces blood loss in cemented hip arthroplasty: a randomized, double-blind study of 39 patients with osteoarthritis. *Acta Orthop*, 2005, 76(6): 829-832.
- [29] Claeys MA, Vermeersch N, Haentjens P. Reduction of blood loss with tranexamic acids in primary total hip replacement surgery. *Acta Chir Belg*, 2007, 107(4): 397-401.
- [30] Rajesparan K, Biant LC, Ahmad M, et al. The effect of an intravenous bolus of tranexamic acid on blood loss in total hip replacement. *J Bone Joint Surg Br*, 2009, 91(6): 776-783.
- [31] Kazemi SM, Mosaffa F, Ejazi A, et al. The effect of

- tranexamic acid on reducing blood loss in cementless total hip arthroplasty under epidural anesthesia. *Orthopedics*, 2010, 33(1): 17.
- [32] Singh J, Ballal MS, Mitchell P, et al. Effects of tranexamic acid on blood loss during total hip arthroplasty. *J Orthop-Surg (Hong Kong)*, 2010, 18(3): 282-286.
- [33] McConnell JS, Shewale S, Munro NA, et al. Reduction of blood loss in primary hip arthroplasty with tranexamic acid or fibrin spray. *Acta Orthop*, 2011, 82 (6): 660-663.
- [34] Rajesh, Malhotra, Vijay Kumar, et al. The use of tranexamic acid to reduce blood loss in primary cementless total hip arthroplasty. *Eur J Orthop Surg Traumatol*, 2011, 21: 101-104.
- [35] Clavé A, Fazilleau F, Dumser D, et al. Efficacy of tranexamic acid on blood loss after primary cementless total hip replacement with rivaroxaban thromboprophylaxis: A case-control study in 70 patients. *Orthop Traumatol Surg Res*, 2012, 98(5): 484-490.
- [36] 傅峥, 张健, 姚海. 氨甲环酸对全髋关节置换术隐性失血的影响. *重庆医科大学学报*, 2012, 37(4): 359-361.
- [37] Imai N, Dohmae Y, Suda K, et al. Tranexamic acid for reduction of blood loss during total hip arthroplasty. *J Arthroplasty*, 2012, 27(10): 1838-1843.
- [38] Alshryda S, Mason J, Sarda P, et al. Topical (intra-articular) tranexamic acid reduces blood loss and transfusion rates following total hip replacement: a randomized controlled trial (TRANX-H). *J Bone Joint Surg Am*, 2013, 95(21): 1969-1974.
- [39] Wang C, Xu GJ, Han Z, et al. Topical application of tranexamic acid in primary total hip arthroplasty: a systematic review and meta-analysis. *Int J Surg*, 2015, 15: 134-139.
- [40] Van Elst C, Vanbervliet J, Simon JP, et al. The effect of topical application of tranexamic acid in total hip arthroplasty through the direct anterior approach. American Academy of Orthopaedic Surgeons Annual Meeting, March, 2013.
- [41] Martin JG, Cassatt KB, Kincaid-Cinnamon KA, et al. Topical administration of tranexamic acid in primary total hip and total knee arthroplasty. *J Arthroplasty*, 2014, 29(5): 889-894.
- [42] Yue C, Kang P, Yang P, et al. Topical application of tranexamic acid in primary total hip arthroplasty: a randomized double-blind controlled trial. *J Arthroplasty*, 2014, 29 (12): 2452-2456.
- [43] 岳辰, 谢锦伟, 蔡东峰, 等. 静脉联合局部应用氨甲环酸减少初次全髋关节置换术围手术期失血的有效性及安全性研究. *中华骨与关节外科杂志*, 2015, 8(1): 44-48.
- [44] Chen X, Zhu X, Yang S, et al. Tranexamic acid treatment decreases hidden blood loss in total knee arthroplasty. *Am J Ther*, 2015 Mar 12, [Epub ahead of print].
- [45] Wu Q, Zhang HA, Liu SL, et al. Is tranexamic acid clinically effective and safe to prevent blood loss in total knee arthroplasty? A meta-analysis of 34 randomized controlled trials. *Eur J Orthop Surg Traumatol*, 2015, 25(3): 525-541.
- [46] Tan J, Chen H, Liu Q, et al. A meta-analysis of the effectiveness and safety of using tranexamic acid in primary unilateral total knee arthroplasty. *J Surg Res*, 2013, 184(2): 880-887.
- [47] Fu DJ, Chen C, Guo L, et al. Use of intravenous tranexamic acid in total knee arthroplasty: a meta-analysis of randomized controlled trials. *Chin J Traumatol*, 2013, 16(2): 67-76.
- [48] Yang ZG, Chen WP, Wu LD. Effectiveness and safety of tranexamic acid in reducing blood loss in total knee arthroplasty: a meta-analysis. *J Bone Joint Surg Am*, 2012, 94 (13): 1153-1159.
- [49] Alshryda S, Sarda P, Sukeik M, et al. Tranexamic acid in total knee replacement: a systematic review and meta-analysis. *J Bone Joint Surg Br*, 2011, 93(12): 1577-1585.
- [50] Zhang H, Chen J, Chen F, et al. The effect of tranexamic acid on blood loss and use of blood products in total knee arthroplasty: a meta-analysis. *Knee Surg Sports Traumatol Arthrosc*, 2012, 20(9): 1742-1752.
- [51] Cid J, Lozano M. Tranexamic acid reduces allogeneic red cell transfusions in patients undergoing total knee arthroplasty: results of a meta-analysis of randomized controlled trials. *Transfusion*, 2005, 45(8): 1302-1307.
- [52] 张阳, 钱齐荣, 吴海山, 等. 氨甲环酸减少全膝关节置换术失血量的Meta分析. *中华骨科杂志*, 2009, 29(6): 524-529.
- [53] 傅德杰, 陈凯宁, 杨柳. 氨甲环酸对全膝关节置换术失血量影响的系统评价. *中国矫形外科杂志*, 2012, 20(13): 1172-1177.
- [54] 方志辉, 杨华清, 李兵奎, 等. 氨甲环酸应用于膝关节置换术随机对照安慰剂试验的Meta分析. *中华临床医师杂志(电子版)*, 2012, 6(24): 8173-8179.
- [55] Benoni G, Fredin H. Fibrinolytic inhibition with tranexamic acid reduces blood loss and blood transfusion after knee arthroplasty: a prospective, randomised, double-blind study of 86 patients. *J Bone Joint Surg Br*, 1996, 78(3): 434-440.
- [56] Camarasa MA, Ollé G, Serra-Prat M, et al. Efficacy of ampicaproc, tranexamic acids in the control of bleeding during total knee replacement: a randomized clinical trial. *Br J Anaesth*, 2006, 96(5): 576-582.
- [57] Dhillon MS, Bali K, Prabhakar S. Tranexamic acid for control of blood loss in bilateral total knee replacement in a single-stage. *Indian J Orthop*, 2011, 45: 148-152.
- [58] Ellis MH, Freedman B, Zohar E, et al. The effect of tourniquet application, tranexamic acid, and desmopressin on the procoagulant and fibrinolytic systems during total knee replacement. *J Clin Anesth*, 2001, 13(7): 509-513.
- [59] Good L, Peterson E, Lisander B. Tranexamic acid decreases external blood loss but not hidden blood loss in total knee replacement. *Br J Anaesth*, 2003, 90(5): 596-599.
- [60] Hiippala S, Strid L, Wennstrand M, et al. Tranexamic acid (Cyklokapron) reduces perioperative blood loss associated with total knee arthroplasty. *Br J Anaesth*, 1995, 74(5): 534-537.
- [61] Hiippala ST, Strid LJ, Wennerstrand MI, et al. Tranexamic acid radically decreases blood loss and transfusions associated with total knee arthroplasty. *Anesth Analg*, 1997, 84: 839-844.
- [62] Jansen AJ, Andreica S, Claeys M, et al. Use of tranexamic acid for an effective blood conservation strategy after total knee arthroplasty. *Br J Anaesth*, 1999, 83(4): 596-601.
- [63] Kakar PN, Gupta N, Govil P, et al. Efficacy and safety of

- tranexamic acid in control of bleeding following TKR: a randomized clinical trial. Indian J Anaesth, 2009, 53(6): 667-671.
- [64] MacGillivray RG, Tarabichi SB, Hawari MF, et al. Tranexamic acid to reduce blood loss after bilateral total knee arthroplasty: a prospective, randomized double blind study. J Arthroplasty, 2011, 26(1): 24-28.
- [65] Orpen NM, Little C, Walker G, et al. Tranexamic acid reduces early post-operative blood loss after total knee arthroplasty: a prospective randomized controlled trial of 29 patients. Knee, 2006, 13(2): 106-110.
- [66] Tanaka N, Sakahashi H, Sato E, et al. Timing of the administration of tranexamic acid for maximum reduction in blood loss in arthroplasty of the knee. J Bone Joint Surg Br, 2001, 83(5): 702-705.
- [67] Veien M, Sørensen JV, Madsen F, et al. Tranexamic acid given intraoperatively reduces blood loss after total knee replacement: a randomized, controlled study. Acta Anaesthesiol Scand, 2002, 46(10): 1206-1211.
- [68] 胡旭栋, 周宗科, 裴福兴, 等. 全膝关节置换围手术期氨甲环酸不同使用方法的有效性和安全性. 中华骨科杂志, 2014, 34(6): 599-604.
- [69] Panteli M, Papakostidis C, Dahabreh Z, et al. Topical tranexamic acid in total knee replacement: a systematic review and meta-analysis. Knee, 2013, 20 (5): 300-309.
- [70] Zhao-yu C, Yan G, Wei C, et al. Reduced blood loss after intra-articular tranexamic acid injection during total knee arthroplasty: a meta-analysis of the literature. Knee Surg Sports Traumatol Arthrosc, 2014, 22(12): 3181-3190.
- [71] Yue C, Pei F, Yang P, et al. Effect of topical tranexamic acid in reducing bleeding and transfusions in TKA. Orthopedics, 2015, 38(5): 315-324.
- [72] 高福强, 孙伟, 郭万首, 等. 局部应用氨甲环酸减少全膝关节置换术后失血量的系统评价. 中国骨与关节损伤杂志, 2014, 29(8): 772-775.
- [73] Alshryda S, Mason J, Vaghela M, et al. Topical (intra-articular) tranexamic acid reduces blood loss and transfusion rates following total knee replacement: a randomized controlled trial(TRANX-K). J Bone Joint Surg Am, 2013, 95(21): 1961-1968.
- [74] Sa-Ngasoongsong P, Wongsak S, Chanplakorn P, et al. Efficacy of low-dose intra-articular tranexamic acid in total knee replacement; a prospective triple-blinded randomized controlled trial. BMC Musculoskelet Disord, 2013, 14: 340.
- [75] Martin JG, Cassatt KB, Kincaid-Cinnamon KA, et al. Topical administration of tranexamic acid in primary total hip and total knee arthroplasty. J Arthroplasty, 2014, 29(5): 889-894.
- [76] Konig G, Hamlin BR, Waters JH. Topical tranexamic acid reduces blood loss and transfusion rates in total hip and total knee arthroplasty. J Arthroplasty, 2013, 28(9): 1473-1476.
- [77] Georgiadis AG, Muh SJ, Silverton CD, et al. A prospective double-blind placebo controlled trial of topical tranexamic acid in total knee arthroplasty. J Arthroplasty, 2013, 28(8 Suppl): 78-82.
- [78] Roy SP, Tanki UF, Dutta A, et al. Efficacy of intra-articular tranexamic acid in blood loss reduction following primary unilateral total knee arthroplasty. Knee Surg Sports Traumatol Arthrosc, 2012, 20(12): 2494-2501.
- [79] Ishida K, Tsumura N, Kitagawa A, et al. Intra-articular injection of tranexamic acid reduces not only blood loss but also knee joint swelling after total knee arthroplasty. Int Orthop, 2011, 35(11): 1639-1645.
- [80] Maniar RN, Kumar G, Singhi T, et al. Most effective regimen of tranexamic acid in knee arthroplasty: a prospective randomized controlled study in 240 patients. Clin Orthop Relat Res, 2012, 470(9): 2605-2612.
- [81] Onodera T, Majima T, Sawaguchi N, et al. Risk of deep venous thrombosis in drain clamping with tranexamic acid and carbazochromesodium sulfonate hydrate in total knee arthroplasty. J Arthroplasty, 2012, 27(1): 105-108.
- [82] Wong J, Abrishami A, El Beheiry H, et al. Topical application of tranexamic acid reduces postoperative blood loss in total knee arthroplasty: a randomized, controlled trial. J Bone Joint Surg Am, 2010, 92(15): 2503-2513.
- [83] Sa-Ngasoongsong P, Channoom T, Kawinwonggwit V, et al. Postoperative blood loss reduction in computer-assisted surgery total knee replacement by low dose intra-articular tranexamic acid injection together with 2-hour clamp drain: a prospective triple-blinded randomized controlled trial. Orthop Rev (Pavia), 2011, 3(2): e12.
- [84] Seo JG, Moon YW, Park SH, et al. The comparative efficacies of intra-articular and IV tranexamic acid for reducing blood loss during total knee arthroplasty. Knee Surg Sports Traumatol Arthrosc, 2013, 21(8): 1869-1874.
- [85] Huang Z, Ma J, Shen B, et al. Combination of intravenous and topical application of tranexamic acid in primary total knee arthroplasty: a prospective randomized controlled trial. J Arthroplasty, 2014, 29(12): 2342-2346.
- [86] 中华医学会骨科分会. 中国骨科大手术静脉血栓栓塞症预防指南. 中华骨科杂志, 2009, 29(6): 602-604.